200300054

No.



TO ALL TO WHOM THESE PRESENTS SHALL COME:

Progeny Advanced Genetics, Inc.

MICCOLF, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLEMISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR MIDORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE 13E, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT VARIETY THEREFROM, TO THE EXTENT THE PLANT VARIETY PROTECTION ACT. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

LETTUCE

'Cyclone'

In Vestimonn Thereof, I have hereunto set my hand and caused the seal of the Hunt Huristy Heaterton Office to be affixed at the City of Washington, D.C. this fourteenth day of February, in the year two thousand and six.

Atlast:

QCM3

Commissioner Plant Variety Protection Office Agricultural Marketing Service Scretary of Agriculture

U.S. DEPARTMENT OF AGRICULTURE The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995. AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2426). APPLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE (Instructions and information collection burden statement on reverse) 1. NAME OF OWNER TEMPORARY DESIGNATION OR EXPERIMENTAL NAME 3. VARIETY NAME PROGENY ADVANCED GENETICS, INC. CYCLONE 4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) 5. TELEPHONE (include area code) FOR OFFICIAL USE ONLY 1536 B MOFFETT ST **PVPO NUMBER** SALINAS CA 93905-3342 831-751-6030 20030005 6. FAX (include area code) 831-751-6032 FILING DATE IF THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) 8. IF INCORPORATED, GIVE STATE OF INCORPORATION 9. DATE OF INCORPORATION 12/9/2002 CORPORATION CALIFORNIA 12/26/94 10. NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THIS APPLICATION. (First person listed will receive all papers) FILING AND EXAMINATION FEES: DARRYN GIBSON \$ 2705.00 MANAGER OF RESEARCH AND DEVELOPMENT PROGENY ADVANCED GENETICS, INC 1536 B MOFFETT ST 5 SALINAS CALIFORNIA 93905 CERTIFICATION FEE: \$ 682.00 11/3/2005 11. TELEPHONE (Include area code) 12. FAX (Include area code) 14. CROP KIND (Common Name) 831-751-6030 831-751-6032 DMAN@IPROGENY.. COM LETTUCE 15. GENUS AND SPECIES NAME OF CROP 16. FAMILY NAME (Botanical) 17. IS THE VARIETY A FIRST GENERATION HYBRIO? LACTUCA SATIVA COMPOSITAE YES M NO 18. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? See Section 83(a) of the Plant Variety Protection Act) а. 💢 Exhibit A. Origin and Breeding History of the Variety YES (If "yes", answer items 20 and 21 below) NO (If "no", go to item 22) Exhibit B. Statement of Distinctness Exhibit C. Objective Description of Variety 20. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF CLASSES? YES **⊠** NO Exhibit D. Additional Description of the Variety (Optional) IF YES, WHICH CLASSES? FOUNDATION REGISTERED CERTIFIED Exhibit E. Statement of the Basis of the Owner's Ownership Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties, verification that tissue culture will be deposited and maintained in an approved public 21. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? M M Filing and Examination Fee (\$2,705), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office) IF YES, SPECIFY THE FOUNDATION REGISTERED CERTIFIED NUMBER 1,2,3, etc. (If additional explanation is necessary, please use the space indicated on the reverse.) HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U. S. OR OTHER COUNTRIES? 23. IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)? YES 12/31/01, U.S.A X YES PLEASE SEE ATTACHED NO IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.) IF YES, PLEASE GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED REFERENCE NUMBER. (Flease use space indicated on reverse.) 24. The owners declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate. The undersigned owner(s) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Owner(s) is(are) informed that false representation herein can jeopardize protection and result in penalties.

	· · · · · · · · · · · · · · · · · · ·
SIGNATURE OF OWNER	SIGNATURE OF OWNER
V Mente	
AME (Please print or type)	NAME (Please print or type)
DADDYN CTDCON	

DARRYN GIBSON

MANAGER OF RESEARCH AND EVELOPMENT

DATE

CAPACITY OR TITLE

DAT

(See reverse for instructions and information colle

01) designed by the Plant Variety Protection Office with WordPerfect 6.0a. Replaces STD-470 (02-99) which is obsolete.

BRUSSELS
CHICAGO
DENVER
DETROIT
JACKSONVILLE
LOS ANGELES
MADISON
MILWAUKEE
ORLANDO
SACRAMENTO
SAN DIEGO/DEL MAR
SAN FRANCISCO
TALLAHASSEE
TAMPA
WASHINGTON, D.C.
WEST PALM BEACH

200300054 FOLEY LARDNER

February 26, 2002

Darryn Gibson Progeny Advanced Genetics 1536 B Moffet Street Salinas, CA 93905

Re:

U.S. Provisional Patent Application No. 60/339,808

Title:

Lettuce Variety PX 408

Inventor(s):

Nathan K. Olivas; Nathan J. Olivas

Filing Date:

December 14, 2001

Our Ref.:

087558-0103

Dear Darryn:

Enclosed is the following in regard to the above-identified application.

The Provisional Filing Receipt:

Serial No.: 60/339,808

Filing Date: December 14, 2001

We will keep you informed of further developments as they occur. If you have any questions regarding this application, please do not hesitate to contact us.

Best regards,

Richard C. Peet

Nina Osseiran

RCP/NMO:ljp Enclosure(s)

FOLEY & LARDNER WASHINGTON HARBOUR 3000 K STREET, N.W., SUITE 500 WASHINGTON, D.C. 20007-5143

TEL: 202.672.5300 FAX: 202.672.5399 WWW.FOLEYLARDNER.COM WRITER'S DIRECT LINE 202.945-6180

EMAIL ADDRESS nosseiran@foleylaw.com

CLIENT/MATTER NUMBER 087558-0103

002.733985.1



Exhibit A

Details of the Development of the Variety Cyclone

Cyclone is a unique romaine lettuce variety developed from a hand pollinated cross of Darkland Cos and Major Cos made in 1993 in the San Joaquin Valley. The F1 seed harvested was designated as Gamma # 035. Darkland Cos, a dark green PIC type available from Central Valley Seed was selected as a source of dark color. Major Cos, developed by Genecorp Seed was a source of weight and adaptability. The cross was made, and by the incorporation of the pedigree selection breeding method we developed a dark green, slow bolting romaine variety.

Approximately 50 plants of the F1 seed were planted in a San Joaquin Valley production field for seed increase in 1994. The F2 seed was harvested in August 1994, labeled 94035.

Line 94035 was planted in 1996 in a San Joaquin Valley research production block. Individual plants were selected at market maturity for distinctions in type, size, color and maturity. The particular selection labeled PSR95400-47 stake # 422-1 was noted to be extremely dark in color, have an unusually thick leaf texture, and be significantly shorter and more compact than the remainder of the plants in the block. The selected plant was allowed to mature and the F3 seed from the selection was harvested individually.

The F3 seed from selection number PSR95400-47 stake #422-1 was increased in our 1997 SJV research production block. The block was evaluated and rogued at market maturity removing any visible off-types. The selected remaining plants exhibiting the dark color, thick leaf texture and compact growth habit were allowed to self pollinate and the F4 seed was harvested in bulk in the fall of 1997. The F4 seed was labeled SJ97-Stake #842C-W.

The F4 seed was evaluated in research and development plot trials during the 1998 growing season in the Salinas Valley of California and Yuma Arizona, where it exhibited excellent uniformity demonstrating the selected traits. Recommendation for increase was made, and the F4 seed was planted in our San Joaquin Valley Research production field in 1999. The block was planted and identified as Stake #99B1550. The line was evaluated and rogued first at market maturity, where it was noted as uniform, demonstrating the desired selected traits. Selective rogueing was done, removing earlier maturing plants. The F5 seed was harvested in bulk.

The F5 line was identified as PX 408 in January of 2000 and was trialed through out the Salinas Valley where it demonstrated excellent uniformity in terms of size, type and maturity. The variety, during this trialing period was identified and preliminarily tested for the developing whole leaf and baby leaf lettuce markets. The F5 seed was then increased in our 2000 San Joaquin Valley commercial production field as Stake #00K574. The variety was noted as uniform and stable without variants. The F6 seed was harvested in the fall of 2000.

During the remainder of 2000 and 2001 further trials and tests were conducted for the romaine whole leaf market, through a patented process. The variety tested extremely well, offering thick heavy leafs uniform in size, shape and color. During this same time period trials for the baby leaf market were also conducted, where it also tested extremely well. Based on this trial performance the F6 seed was again increased in our 2001 production and the F7 seed was harvested in September of 2001. Cyclone as observed in multiple field trials and in our seed production crops, has been uniform, stable, and with out variants for 3 generations. Seed from this variety was first sold on December 31, 2001, no harvested material from this variety was sold prior to this date.



200300054

Exhibit B Statement of Distinctness

Cyclone is a new and distinct variety of romaine lettuce, that most closely resembles Darkland Cos. This variety is distinguished by its distinct plant height, which is shorter than typical romaine. Cyclone is also distinct as it produces thicker leafs with a darker color, and a more uniform size than other romaine varieties.

The following characteristics distinguish Cyclone from its parent variety Major Cos:*

- 1. Cyclone is darker in color than Major Cos. Cyclone measured 7.5 gy4/6 and 7.5 gy4/4 versus Major cos with a color rating of 5gy 4/6 and 5gy 4/8
- 2. Cyclone averages 4 days later maturing than Major Cos
- 3. Cyclone has a statistically significant shorter heart length
- 4. Cyclone has a statistically significant shorter head length
- 5. Cyclone has a statistically significant smaller frame diameter
- 6. Cyclone has a statistically significant shorter leaf
- 7. Cyclone has a statistically significant thicker leaf

Additional distinctions as cited in their respective Exhibit C documentation:

Characteristic	Cyclone	Major Cos
Cotyledon shape	Spatulate	Intermediate
Cotyledon leaf undulation	Flat	Medium
4 th leaf reflexing	None	Lateral margin
Leaf thickness	Thick	Intermediate
Bolter leaves	Curved	Straight
Bolter leaf margins	Dentate	Entire

The following characteristics distinguish Cyclone from its parent variety Darkland Cos:*

- 1. Cyclone is darker in color than Darkland Cos. Cyclone measured 7.5 gy4/6 and 7.5 gy4/4 versus Darkland Cos with a color rating of 5gy 5/6 and 5gy 4/8.
- 2. Cyclone is typically 2-6 days later maturing than Darkland Cos.
- 3. Cyclone has a statistically significant shorter head length
- 4. Cyclone has a statistically significant smaller frame diameter
- 5. Cyclone has a statistically significant shorter leaf
- 6. Cyclone has a statistically significant thicker leaf





Additional distinctions as cited in their respective Exhibit C documentation:

Characteristic	Cyclone	Darkland C
Cotyledon shape	Spatulate	Broad
Cotyledon leaf shape	Elongated	Oval
Anthocyanin distribution	Absent	Throughout
Cotyledon leaf cupping	Slight	Uncupped
Margin undulation	Moderate	Absent
But midrib	Moderate raised	Flat
Bolter leaves	Curved	Straight
Bolter leaf margins	Dentate	Entire
Bolter terminal inflorescence	Present	Absent
Bolter basal side shoots	Absent	Present

The following characteristics distinguish Cyclone from the variety King Henry:

- 1. Cyclone has a smother leaf surface than the variety King Henry. The leaf surface of King Henry is described as heavily savoyed or blistered, where as the leaf surface of Cyclone is described as moderately savoyed or blistered.
- 2. Cyclone is darker in color than King Henry. Cyclone measured 7.5gy4/6 and 7.5gy4/4 versus King Henry with a color rating of 5gy 4/6.
- 3. Cyclone is a shorter growing plant than King Henry. On average King Henry averages a plant height of 270 cm's. On average, Cyclone averages a plant height of 251 cm's.

The following characteristics distinguish Cyclone from the variety Frontier Cos:

- 1. Cyclone has a more savoyed leaf surface than Frontier Cos. The leaf surface of Frontier is described as smooth, where as the leaf surface of Cyclone is described as moderately savoyed or blistered.
- 2. Cyclone is darker in color than Frontier. Cyclone measured 7.5gy4/6 and 7.5gy4/4 versus Frontier with a color rating of 5gy5/6.
- 3. Cyclone produces a higher yield of useable leaf product than the variety Frontier. On average Cyclone will produce 1.61 finished 10 pound cartons of leaf product from a single tray of 24 plants. Frontier on average produces on average 1.11 finished 10 pound cartons of leaf product from a single tray of 24 plants.

These characteristics are distinct and offer a significant advantage for whole leaf processing by increasing the percent recovery during the patented post harvest process. These characteristics also offer an advantage for the baby leaf lettuce market.

The leaf thickness prevents mechanical damage that can occur during the leaf removal and washing processes, increasing the percentage useable product from the plant.

^{*}All data is supported statistically as seen in the following tables of trial results conducted in the Salinas Valley in the 2001 and 2002 growing seasons.



The whole leaf market specifies strict leaf sizing for acceptable product. The leaf size of a typical romaine is larger than what is acceptable in the whole leaf market. To get leaves of acceptable sizing, the larger outer leaves and the smaller inner leaves must be discarded. Due to the short growth habit, and uniform leaf size of Cyclone, this variety produces a high number of leaves that fall with in this specified range of acceptance. This distinction is of economic benefit as more of the product from each plant can be processed.

U. S. Department of Agriculture Agricultural Marketing Service Science and Technology Program

OBJECTIVE DESCRIPTION OF VARIETY LETTUCE Lactuca sativa

NAME OF APPLICANT (S) FOR OFFICIAL USE ONLY PROGENY ADVANCED GENETICS, INC. ADDRESS (Street and No. or R.F.D. No., City, State, and ZIP Code) 1536 B MOFFETT STREET SALINAS CALIFORNIA 93905 EXPERIMENTAL DESIGNATION Place numbers in the boxes for the characters which best describe this variety. Measured data should be the mean of an appropriate number (at least 10) of v spaced plants. Royal Horticultural Society or any recognized color standard may be used to determine plant colors. The location of the test area is: Color System Used: UNSEL (See list of suggested check verieties page 4.) 01-Curting/Leaf 05-Great Lakes Group 09-Stem 02-Butterhead 06~Vanguard Group 10-Latin 03~Bibb 07=Imperial Group 11-OTHER-04-Cos or Romaine 08*Eastern (Ithaca) Group COLOR HEAT DORMANCY LIGHT DORMANCY 2. SEED: 1-White (Silver Gray) 1-Light Required 1=Susceptible 2-Black (Gray Brown) 2-Not Süsceptible 3-Brown (Amber) 2-Light Not Required 3. COTYLEDON TO FOURTH LEAF STAGE: NOTE: Provide a color photograph or photocopy of the fourth leaf from 20 day old seedling grown under optimal conditions. SHAPE OF COTYLEDONS: 1~Broad 2-Intermediate 3-Spatulate SHAPE OF FOURTH LEAF: 2 LENGTH/WIDTH INDEX OF FOURTH LEAF: L/W x 10 APICAL MARGIN: 1-Entire 4-Moderately Dentate 7-Lobed 2-Creanate/Gnawed 5-Coarsely Dentate 8-OTHER (specify) BASAL MARGIN: 3-Finely Dentate 6-Incised 4=Marked UNDULATION: 1=Flat 2-Slight 3-Medium 7-Gray Green 3-Medium Green 1-Yellow Green 5-Blue Green GREEN COLOR: 2=Light Green 4-Dark Green 6-Silver Green ANTHOCYANIN:

3-Spotted

2-Moderate

2-Present

2-Slight

2-Apical Margin

4-Throughout

1-Absent

1-Light

1-Absent

1-None

1-Uncupped

2-Margin Only

DISTRIBUTION:

ROLLING:

CUPPING:

REFLEXING:

CONCENTRATION:

5-OTHER (specify)

3-Intense

3-Markedly

3-Lateral Margins

		LS (boserve harvest-		, '	
. NC	OTE: P	rovide color photo of herve MARGIN:	st-mature leaves which accurately shows color	and mergin characteristics.	
		INCISION DEPTH: Ideepest penetration	1=Absent/Shaflow (Dark Green Boston of the margin)	n] 2~Moderate {Vanguard}	3-Deep (Great Lakes 6)
		INDENTATION: (finest divisions of the ma	1=Entire (Dark Green Boston) 2=Shallowly Dentate (Great Lakes 65)	3-Despty Dentate (Great Lakes 659 4-Crenate (Vanguard)	1 S-OTHER (apecify)
	2	UNDULATION OF APICAL MARGIN:	THE 1+Absent/Slight (Dark Green Boston)	2-Moderate (Vanguard)	3-Strong (Great Lakes 6
	5	GREEN COLOR:	1-Very Light Green (Bibb) 2-Light Green (Minetto)	3-Medium Green (Great Lakes) 4-Dark Green (Vanguard)	5-Very Dark Green 6-OTHER
		ANTHOCYANIN (grown	at or below 10 C):		
		DISTRIBUTION:	1-Absent 2-Margin Only (Big Boston)	3-Spotted (Calif, Cream Butter) 4-Throughout (Prize Head)	5-OTHER (apecify)
		CONCENTRATION:	1=Light (Iceberg)	2-Moderate (Prize Head)	3-Intense (Ruby)
	2	SIZE:	1-Small	2-Medium	3=Large
· .	2	GLOSSINESS:	1=Dull (Variguard)	2-Moderate (Selinas)	3-Glossy (Great Lakes)
	2	BLISTERING:	1=Absent/Slight (Satinas)	2=Moderate (Vanguard)	3=Strong (Prize Head)
	3	LEAF THICKNESS:	1-Thia	2=Intermediate	3=Thick
		TRICHOMES:	1=Absent (smooth)	2-Present (spiny)	
5. PLAN	IT (at m	arket stage. Choose a comp	parison variety appropriate for this type.):		
3	4 s	This Variety	ES: DARKLAND	COS (specify comparison variety	r)
3		HEAD DIAMETER (market m This Variety	trimmed with single cap leaf): 38 cm DARKLAND	COS (specify comparison variety	-)
[4	EAD SHAPE:	1-Flattened 2-Slightly Flattened		5=Non-Heading 6=OTHER
		EAD SIZE CLASS:	1-Small	2-Medium	3=Large
2	<u> </u>	EAD COUNT PER CARTO	N		
77	<i>(</i>) (EAD WEIGHT: This Variety	1863. DAPKLAND C	OS (specify comparison variety,	
	=	EAD FIRMNESS:	1=Loose 2=Moderate	3–Firm 4–Very Firm	
6. BUTT	bottom	of market-trimmed head):			
	3]_sh	IAPE:	1-Stightly Concave	2~Flat 3	-Rounded
7 5005 (DRIB:	1-Flattened (Salinas)	2-Moderately Raised 3	=Prominently Raised (Grea Lakes 659)
39	57	market-trimmed head): Diameter at base of head	,	-	
181	7 Rai	io of head diameter/core di	ameter		
49	<i>,</i> ,	e height from base of head: This Variety	to apex: 49 mm DARKLAND CO	se.	
8. BOLTING				Ispecify comparison variety	UTE
62		nber of days from First Wat Variety	er Date to seed stalk emergence (summer co	nditlone);	
12	<u> </u>	TING CLASS-	7_C1+		Very Rapid
1118		ht of mature seed stalk: This Variety	MADRIAND C	-Rapid	<u>a</u>
	ا لي		TIPINION TOURNING (OS (specify comparison variety)	• •

	• ` .			200300054
29	Spread of Bolter Plant fot u	ildest point): 3 @ cm .	VIARVIAND CAC	'y comparison variety)
2	BOLTER LEAVES:	1=Straight	2-Curved	,
2	MARGIN:	1=Entire	2+Dentate	
[3]	COLOR:	7-Light Green	2-Medium Green	3-Dark Green
2	BOLTER HABIT: TERMINAL INFLORESCENCE:	1-Absent	2-Present	
2	LATERAL SHOOTS:	1-Absent	2-Present	
	BASAL SIDE SHOOTS:	1=Absent	2-Present	
9. MATURITY	carliness of harvest-mature h	ead formation):		
NOTE: Co	omplets this section for at les	t one season.		
SEASC	N Applic 1 #of days	Check 1/ #of days	- CHECK	C VARIETY 3
Spring	89	85	DARKLAND COS	
Summer	66	62	DARKLAND COS	
Fall	70	68	DARKLAND COS	
Winter	98	92	DARKLAND COS	
Give planting da	te(s), and location(s):			
Spring	MARCH 10, S	VINAS VALL	EY CA	
Summer	JUNE 27, SA	LINAS VALLE	EY CA	
			EYCA; SEPT 22,	YUMA AZ
Winter FEB 4 SALINAS VALLEY CA 1/ First water date to harvest. 2/ Fill in check variety name on the appropriate line.				
2) First water date to harvest. 2) Fill in check variety name on the appropriate line. ADAPTATION:				
PRIMARY REGIONS OF ADAPTION (tested and proven adapted): (0~Not tested 1=Not Adapted 2=Adapted)				
2 Sou	thwest (Colif., Ariz. desert)	Z West Coas	st O Northeast	
O Nort	thcentral	O Southeast		· · · · · · · · · · · · · · · · · · ·
SEA	SON: Spring larea <u>SAUNAS V</u>	ALLEV .	Z Fall (areaSALNAS)	VALLEY, YUMA
	Summer (area SALINAS	1	Winter (area SALM)	
O GAE	ENHOUSE: 0-	Not tested	1=Not Adapted	2-Adapted
3 soil	TYPE: 1-A	1ineral	2-Organic	3-Both

Page 3 of

FORM LS-470-1 (9-86)

200300054

11. DISEASES ANI	D STRESS REACTIONS	(O-Not tested: 1-	·Suscaotibie: Z=In	termediate: 3=Resist	cant: 4-Highly resistant: 5-Tolerant):
	VIAUS				FUNGAL/BACTERIAL
	Big Vein			Carky Root R	ot (Pythium Root Rot)
	Lattuca Mosaic			2 Downy Milder	~ (Reces)
	Oucumber Mosaic	•		5 Powdery Milde	ew
	Sroad Sean Wilt			2 Scieratina Ra	en de la companya de La companya de la co
	O Turnia Massic			3 Bacterial Soft	Rot (Pseudomonas spa. & others)
•	Beet Western Yellov	vt	4	3 Sotrytis (Gray	
	Latt. Infectious Yes	laws		OTHER	
	Other Virus				
		<u> </u>	············	•	
	INSECTS			•	PHYSIOLOGICALISTRESS
	Cabbage Loopers		•	5 Tracoum	O Sait
	O Root Achids			5 Heat	Srown Rib (Rib Discoloration, Rib Stign
	OGreen Peach Achid			Oraught	OTHER
	Other Insect			2 Caid	
	- 		_ · ·		
•			POST HARVES		
	O Plak Rib			5 Internal Rib N	lecrosis (Blackheart, Gray Rib, Gray Streak)
	S Russet Sporting			S Brown Stain	•
	Surv Brown Discol	oration			
12 BICCHEMICAL	OR ELECTROPHORETI	C MARKERS:			
	,				
		•			
13. COMMENTS:			· · · · · · · · · · · · · · · · · · ·		
		•	1 1		

SELTELE AND MORE SELECTION OF TREBUDUS


TYPE


- CUTTING/LEAF BUTTERHEAD


- 51
- BUTTERHEAD BIBB COS. OR ROMAINE GREAT LAKES GROUP VANGUARD GROUP IMPERIAL GROUP EASTERN GROUP STEM LATIN 51 71
- 31
- 91

CHECK VARIETY

SALAD BOWL
DARK GREEN BOSTON
BIBB
PARRIS ISLAND
GREAT LAKES 559-700
VANGUARD AVIV THACA CELTUCE MATCHLESS







Cyclone Leafs





Exhibit D:

Additional Variety Description:

Cyclone is a new and unique variety of romaine lettuce due to its dark color, and growth habit. The growth habit is distinct as it is slower growing and shorter in height than other romaine varieties. Cyclone is also distinguished by the uniformity, texture, and size of its leaves. This variety produces a higher than normal percentage of leaves of the same size. This is a benefit in the whole leaf pack, as more of the leaves from each plant can be used.









Cyclone

Date Mature Days	9/1/01
<u>.</u>	Cyclone
8/28/01	dg ac
Date evald:	Eval by:
5-6 north	al Va PIC
Ranch/lot;	king Commerci
Salinas	Royal Pa
Location:	Grower:
PDS01027	6/27/0
Trial map #:	Wet Date:

,	Date Mature	Days to Maturity	Color
Cyclone	10/1/6	99	7.5gv4//4
Major Cos	10/8/6	89	5gy 4/8

	Core Diameter (mm	r (mm)	Cores Louise Com	(mm	Control of the second	A Company of the Comp	STATE OF THE PERSON NAMED IN COLUMN	220	Section and expenses and expenses					
17 T					TO THE PROPERTY OF THE PARTY OF	· /	I DROLL	Tengra (mm.)	Head length: Core Length		Frame diam (cm)		Head wt. (g)	
Sample #	Cyclone	Major Cos	Cyclone	Major Cos	Cyclone	Major Cos	Cyclone [1	Major Cos	Cyclone	Maior Cos	Cyclone	oior Coe	Cyclone	Marian
_	35	43	35	43	081	210	0.50	1000	1	Ş	1	T	1	Iviajui Cus
2	40	40	**				200	7.70	7,	0.0	\$	30	685	549
					180	200	210	260	3.8	6.5	30	41	806	828
~		45	49	45	180	230	230	290	4.7	4.9	30	38	202	
4	40	52	26	52	091	240	210	300	2,5	0 7	000	\$	201	13.0
43	39	43	54	43	190	740	030	310	7 7	200	07	3 3	/5/	655
°	5 42	45	\$9			02.0	730	010	0.4	7./	OS S	30	830	497
	40	4b	41			2 2	200	ALC	2		75	38	892	721
					,	OH/	240	310	4.7	7.8	31	39	456	575
~	4	39	26	39	170	180	210	280	3.8	7.2	33	Ą	210	089
5	42	42	40	42	160	240	200	320	Ş	76	3	-	717	600
10	40	40	47			4,5		777	A.C	0.7	22	7	/0/	641
11						017	007	300	0.0	7.5	28	4	714	799
	44.1	39				230	250	320	5.1	8.2	32	30	461	475
12	40	45	49	45	170	210	220	290	4.5	6.4	22	33	640	644
Average	39.9	42.8	47.0	42.8	182.5	225.0	229.2	208.2	47	0.4	310	20.1	200	9 900
Stan dev	1,83E+00	3.72E+00	7.29E+00	3.72E+00	1 71E+01	2 30F±01	1 88E±03	1 755 101	1 000 100	2000	0.15	1,75	67/60	630.8
T thet	CO 2777 C	L	O AMES ON		100	100	YA 300	1 /JE ru	1,00ET00	0.602-01	1,935+00	1.44E+00	1.30E+02	1.13E+02
1641 7	4.14E-U2		8.4/E-US		3.Z3E-05		4.22E-09		1.56E-06		8 58E-11		1 0KB 01	

Data highlighted in yellow indicates a statistically significant difference at a minimum 95% confidence level.

ti.	
	Level %
Core Diameter	26
Core length	66
Heart Longth	66
Head length	66
Head length:Core Length	66
Frame diam	66
Leaf Length	66

	Lear Count		Leaf Length		Leaf Width		
	Major Cos	Cyclone	Major Cos	Cyclone	Major Cos	Cyclone	
1	20	22	22	16			
2	21	20	20	16	6	13	
3	19	29	20	10	11	13	
4	19	19	24	19	13	13	
'	22	22	25	16	13	15	
9	21	19	ST.	20	15	13	
7	21	61	17	20	15	15	
8	20	61	87	20	17	17	
φ.	20	21	30	21	14	15	
10	21	20	31		17	18	
11	20	20	31	20	19	18	
12	20	20	33		19	19	
13			16	17	20	15	
14			08	71	20	15	
15			61	17	19	16	
, 16			21	21	21	17	
17			17	20	22	17	
18			20	20	77	14	
19			77	61	20	18	
20			29	10	11	18	Leaf leng
Average	20	21	56	10	41	16	
dard Dev	0.887625365	2.79067712	4.872047014	4.755442664	4.070432539	2.036508881	
T-Test	T-Test 0.560242129		11931 THE -05		0.334247376		

Color	7.5gy4/4
s to Maturity	70
Date Mature av	11/2/01
	Cyclone
10/30/01	CODG
Date evald:	10 Eval by: A
Turry	
Ranch/lot:	Commercial Van
Salinas	Costa/Mann
1038 Location:	8/24/01 Grower:
Trial map #: PDS0	Wet Date:

						~	-				_						.,	_	_	_	
				_		Carolone	a and	7/4					302	401				454	Α.	2.87	10.07100
	Color	7.5gv4/4	5gy5/6		Head w((p)	Major Cos	043	617	110	250	710	617	043	023	704	635	497	965	645 8	6 724-01	1.39E-07
						clone	23	35	36	35	36	6	36	30	3,5	34	35	35	34.8	1 115+00	
	ays to Maturity	۶	<i>L</i> 9		Frame diam (om	Major Cos	ę	25	8	30	00	7	+ +	Ç.	43	38	42.	38	39.0	2 718+00	4.81E-06
	Date Mature	12	10/30/01			Cvelone	77	9	6.9	7.5	1.9	6.3	8 9	7.2	9.9	9,8	7.1	7.2	7.1	6.79F-01	
		Сусіопе	Major Cos		Head length:Core Length	Major Cos	7.7		8.3					5 8			8.3	7.1	7.3	9.38E-01	3.80E-01
					Head length (mm)	Cyclone	270	270	270	270	2.70	0.70	280	280	270	250	290	280	272.5	9.65E+00	
	10/30/01	AC/DG			Head le	Major Cos Cyclone	340						L			320	340		333.3	1.4	2.78E-11
;	Date evald:	10 Eval by:			(0	Cyclone	210	210	220	230	210	230	200	220	230	230	240	220	220,8	1.16E+01	
	Lumy	10			Heart I स्वाहार्क (स्त्रात)	Major Cos	260	300	280	240	270	270	270	260	260	250	260	250	264.2	1.56E+01	1.11E-07
	Kanch/lot:	Commercial Var:				Сусіопе	35	41	39	36	4	43	41	39	41	29	41	39	39.0	4.07E+00	
2.1	Samues	Costa/Mann			Core Length (ann)	Major Cos	44	19	4	35	47	51	49	40	48	49	41	48	46.1	6.73E+00	5.00E-03
Tanadina	ای	8/24/01 Grower:				Сусюпе	34	35	36	37	38	38	39	37	35	36	35	34	36.2	1.64E+00	
01029	Į	8/24/01			Diameter (mm)	r Cos	39	9	41	40	39	41	42	9	39	38	\$	41	40.0	13E+00	.06E-06

Data highlighted in yellow indicates a statistically significant difference at a minimum 95% confidence level

Statistically Distinct Confidence	Trait Level %	Core Diameter 99	Core length 99	Heart Length 99	Head length 99	Frame diam 99	Head Weight 99	Leaf Length 99										16 Leaf length and leaf width numbers are measurements of the useable leafs from one plant.				
12	12	13	16	14	91	17	18	16	17	17	15	19	17	16	17	16	15	161	16	2.01246118		
11	10	13	15	12	13	18	14	16	17	17	20	21	18	19	18	81	19	18	16	3.375998878 2.	0.735138583	
	23 To	25 21		24 21	25	2.0	26 20	28 22	2.5	20 20	60 00	29 21	28 21	29 20	28 19	30 20	71 17	61 10	26 20	8768 1.637552731	E-08	
07	20	17	17	18	19	17	20	20	19	19									19	1.53741223 3.341918768	S183139E-08	
7	17	20	21	24	18	20	21	24	17]	20									20	2.503028469	0.214725829	
7	m	4	50	9		000	Φ,	10		12	13	47	15	16	17	18	19	20	Average		T.Test	

_	_
5/7/02	
Date evald:	of Ferol bay dofar
Salmina 2	Costa/Mann Commercial Va El Camino Real
Ranch/lot:	Commercial
Soledad	Costa/Mann
Location:	2 Grower:
RSV02026	2/4/02 Grower:
#	Wet Date:

o Maturity Color	8 7.5gv4/6	2 5gy4/6
Date Mature Days t	5/13/02	8/7/02
	yclone	Major Cos

	Core Diameter (mm	*(mm)	Core Length (mm	mm)	Heart Length (mm)	(m)	Class langet (rem)	Table Service	TI SAN STATE OF THE PARTY OF TH	Section Community	A STATE OF THE PARTY OF THE PAR			
Sample #	Maior Cos	Cyclone	Major Cos	Crefons	V		PI.	(S)(S)(S)(S)(S)	1		rrame dam (cm)		Head wt. (g)	
			and or cas			Cyclone	Major Cos	Cyclone	Major Cos C	Cyclotte	Major Cos	Cyclone	Major Cos	Cyclone
	51	39	55	09	250	012 210	280	270	4.1	4.5	- 4	ì	Š	
2 4	52	55	53.	20	240	200	200	030	2.2	F	2	Ť,	801	84/
, ,	3	52	**					707	7.6	0.0	4	37	1034	883
							330	280	6.0	4.6	4	32	788	777
			3	99	270	190	340	270	5.2	4.5	157	37	030	003
1	55	39	75	19	270	007	350	270	4.7	44	44	76	1004	200
)	6 52	52	99	62	250	022	320	760	40	4.0	ş	100	1000	£
	7 52	37	9	6.5				2	È	7.4		33	7801	792
3				CO			æ.	720	5.7	3.8	43	30	1219	751
	24	39	70	55	270	190	350	250	5.0	4.5	44	30	816	0.57
-	58	36	45	5.5	280	2000	200	020				3	OFO	100
10	5						one	707	0.7	4.7	4.5	38	1085	982
		1		00	300	210	360	280	5.5	4.7	48	36	1175	1134
Ī	56	36	70	65	280	210	340	260	4.9	4.0	43	38	OEO	3001
12	2 41	55	50	62	290	220	380	270	97	4.4	200	36	0/6	1030
Average	52.6	43.4	2.09	59.7	277.5	ŕ	227.6	264.0			7	CC	775	466
Ston dar	A 36T	1	3	200			536.7	7.4-07	0.0	4.4	44.2	35.4	997.2	915.1
Oran nov	4,6,5,00	WHEN'	9.14E+00	4.36E+00	1.91E+0	1.07E+01	2.83E+01	1.08E+01	8.50E-01	3.17E-01	2.76E+00	2.64F+00	1 47E+02	1 14F+02
T test	1.44E-03		7.35E-01		5,91E-10		8 86F-08		2 802-04		2012 00	20.171.00	10000	70 7117
									1		0/10/10		- Y H	

Indicates that the varieties are significantly different for this trait at least a 95% confidence level

Statistically Distinct	Confidence
Trait	Level %
Core Diameter	9'86
Heart Length	66
Head length	66
Head length: Core Length	66
Frame diam	88
Leaf Count/ Plant	66
Leaf Length	66
LeafWidth	66
Leaf Thickness	66

UB PLABING STREET		The second second second	Constituted and the constitution of the consti	D. S. C.		LOGI THIONIESS (THE	
Major Cos Cyclone	Cyclone	Major Cos Cyclone	Cyclone	Major Cos	Cyclone	Major Cos Cyclone	Cyclone
1 45	35	251	231	001	141	0.93	56.0
	32	272	192	142	126	160	1.22
3 39		292	221	145	136	78.0	260
36		283	245	146	154	0.72	66'0
5 42	26	321	198	951	110	97.0	0.93
41	39	167	233	751	182	08'0	66.0
7 38	37	320	218	781	152	0.70	560
37	35	331	227	177	144	86'0	86'0
4	34	305	242	961	180	98.0	96'0
10 44	34	33I	243	211	155	160	0.82
	32	310	248	185	180	0.85	
12 43	32	331	254	219	187	96'0	66'0
13		336	760	210	177	0.92	0.82
14		320	218	[8]	141	16.0	0.82
15		320	237	061	152	0.75	660
16		314	239	200	081	0.74	60
17		293	252	210	178	6.85	1.4
18		291	248	200	159	26'0	1.29
19		306		202	162	160	96'0
20		313	233	194	160	\$8.0	160
Average 40	33	307	234	781	158	98'0	1.0
	3,84550111		17.79554941	31.8927479	31.8927479 20.94252787	0.087507894	0.158426473
T-Test 1.86E-04		6.80E-14		50°390'L		CHATCH I	

The Contract

RSV0.	RSV02049 Location:		Salinas	Ranch/lot:	1219	Date evald:	5/31/02			Date Mature	Days to Maturity		Color	
	3/10/02 Grower:	ower:	D'arrigo	Commercial Var:	Frontier	Eval by:	AC		Cyclone	6/1/02	68		7.5gy4/4	
	ě								Major Cos	6/2/02	84	7,	5gy4/8	
Core D	Core Diameter (mm)		Core Length (mm)	(mm)	Heart Length (mm	(mn)	Head leneth (mm		Head length Core Length	ore Length	Hrame diam (cm)		Head ort (a)	
Major Cos	П	PX 408	Major Cos	Cyclone	Major Cos	Cyclone	Major Cos Cyclone		Major Cos Cyclone	Cyclone	Major Cos Cyclone	clone		Cyclone
	4	40	50	0 42	235	.5	305	237	6.1	5.6	39	. 35	182	925
27	9	36	3	4 59	9 240	190	341	262	5.3	4.4	35	32	953	979
	\$	41	65	5 47	7 243	3 162	128	229	5.2	4.9	39	33	1136	955
4	\$	37	57	7 57	7 224	4 182	312	239	5.5	4.2	35	34	975	1120
2	39	37	62	2 59.	9 255	5 179	319	252	5.1	4.3	41	33	853	875
9	37	36	49	9 59	9 205	5 192	7.2	269	5.7	4.6	38	32	948	805
<u> </u>	54	35	37	7 62	226	6 205	787	262	9.7	4.2	38	32	1191	1040
œ l	8	4	64	4 57	7 226	6 179	322	239	5.0	4.2	34	34	702	786
6	37	32	51	1 39	216	6 172	277	249	5.4	6.4	38	38	758	995
2	35	45	29	7	4 209	9 159	580	237	4.5	5.4	38	33	885	1013
_	37	9	54	46	5 216	6 166	294	224	5.4	4.9	35	34	698	1156
12	39	42.	67	7 44	1226	691 9	304	205	4.5	4.7	39	34	1004	1286
	39.7	38.4	27.3	3 51.3	3 226.8	8 176.7	8'50E	242.0	5.5	4.8	37.4	33.7	954.7	994.6
3.	3.06E+00	3.55E+00	9.27E+00	0 8.25E+00	1.46E+01	1 1.39E+01	2.15E+01	1:81E+01	8.17E-01	6.83E-01	2.15E+00	1.67E+00	1.56E+02	1.44E+02
3,	3.66E-01		1.08E-0		1:64E-08	95	8.11E-08		4.82E-02		9.21E-05		\$ 22E-01	

2656
100
320
1900
200
35
Ε
×o.
200
.5
A
2
22
<u>o</u>
94
2
a
曼
ŭ
- 3
. 60 - −
- 4 S
o _0
a n
. Q
2 5
<u> </u>
35 45
2 3
, o
4 %
毒 %
13 c
2 6
8 2
= 5
3 -
at at
图 #
and and

Confidence Level %

Statistically Distinct Trait

99 99 95.2 97.5 99 96.8

Heart Length
Head length: Core Length
Franc dian
Leaf Count' Plant
Leaf Length
Leaf Width
Leaf Thickness

	Loaf Comer Plant	lant	Leaf Length		Leaf Width		Leaf Thickness	
Sample #	Major Cos	Cyclone	Major Cos Cyclone	Samblebie	Major Cos	Cyclone	Major Cos	Cyclone
1	37	7	274	177	132	168	0.50	08.0
2	37	7	276	214	142	192	09'0	08'0
3	35	5 36	271	220	061	125	0.50	08'0
4	32	31	254	222	176	186	0.50	09'0
5	26	5 35	271	219	581	177	0.40	08'0
9	32	27	236	582	841	146	0.50	050
7	35	35	237	243	142	152	0.40	0.50
8	34	1 31	241	234	132	144	06:0	09'0
6	39	67 29	297	238	163	164	02:0	02:0
10	38	36	255	232	154	168	09'0	0.50
11	017	0.29	263		174	136	0,40	07.0
12	37	7 33	228	204	149	189	0.50	0.50
13			597	223	152	186	09'0	09'0
14			270	219	162	165	050	05'0
15			252	223	651	139	050	05'0
16			266	214	166	142	05.0	
17			258	622	148	160	050	05'0
18			256	209	145	691	09'0	09'0
19			622	228	681	163	09'0	08'0
20			259	238	192	159	0.50	08'0
Average	35	18 31	257	225	150	162	6.54	910
Standard Dev		3.8336627 3.776923552	15 521(2313	10.88166009	17875526 11 8955018551	18.92227871	0.114248114	0.130182059
T-Test	2,4615.02	10	4.83E-09		3.25E-02		70-H95-7	

			ı			. 1	1008	529	666	825	913	500	4	848	880	862	849	243	213	y IO	প্	7
_					Darkland Cos	20 mmmm /	10	5	6	00	6	Ē	AT.	10	8	8	ŏ	8	•	y	896.5	1.45E+02
Color	7.5gv4//4	5gy 4/8		Head wt. (g)		307			702	737	830	802			719	707	714	194		600	697.3	1,30E+02
Days to Maturity	99	2			Darkland Cos	2.0	36	39	39	40	38	36		9	39	40	35	36	30	300	38.1	1.73E+00
Days to 1	9	62		Frame diam (cm)	Cyclone II	33	36	08	62	28	30	32	1	7 2	55	33	28	32	33	200	31.0	1,95E+00
Date Mature	10/1/61	8/28/01	ſ		Darkland Cos	ос У		4.7	770	7.3	4.9	5.4	24	i,	0,1	5.3	5.6	5.3	5.2	2.2	0.0	7.78E-01
	Cyclone	Darkland Cos	770.41. 4 0	riead length: Core Length		5.6	\$ 25	2.5	1.0	5.3	5.0	4.6	4.0	0 7		277	4.5	4.5	5.1	0.5	2 757	3.72E-01
			Heed loant ((mm)	Darkland Cos Cyclone	290	280	090	202	200	290	270	260	200	900	310	310	290	270	285.0	1 TOE-01	1./36/1
8/28/01	dg ac		LTegal for		Cyclone	250	210					230	240						220	229.2	٦	1 490 07
Date evald:	Eval by:				Darkland Cos	210	200	190	010	017	027	190	13	220	320	777	040	210	190	207.5	1 60B+01	TANK TANK
5-6 north			Heart Loneth (mm)	1	1000	180	180	180	160	907	061	180	200	170	160	01.0	0.77	OI7	170	182.5	1.71E+01	1.27E-03
Ranch/lot: 5	1		1	Darkland Con	ľ	90	09	50	40	9	6	nc ;	39	65	50	**	3	000	22	52.8	7.78E+00	
Salinas I Roval Packino			Core Longth (mm)		1		40	45	40	Ş			49	43	35	55				45.8	6.15E+00	2.29E-02
27 Location: 6/27/01 Grower:			(cau	Darkland Cos Cyclone				42	42	43				42	46	43				42.1	2.02E+00	
<u>%</u>			Core Diameter (mm)	Cyclone	35		3	39	40	39	42	Ę		41	42	4	41			39,9	1.83E+00	1.16E-02
Trial map #: PDS01027 Wet Date: 6/2				Sample #		,	7	3	4	5	9	7		*	6	10	11	12	ı	Average	Stan dev	T test

ighlighted in yellow indicates a statistically significant difference at a minimum 95% confidence level Statistically Distinct Confidence Level % Core Dismeter 99 Core length 98 Heart Length 99
--

	Leaf Count/Plant	1t	Leaf Length		Leaf Width		
	Darkland Cos	Cyclone	Darkland Che	Continue	Dorbland Oce	Content	;
_				Amoralia	Carriagu Cos	Сустопе	
				91	9	. 13	
7	20	20	21	91	6	13	<u> </u>
m	20		25	11	12		
4	19	19	24	61			
*	20	22		16			
9	21	61		20		13	
7	22	61				15	
90	22	19		20	18	17	
O.	19	21			16	15	
10	22	20	28			18	
11	22	20	29	20		18	
12	22	20		21	17	19	
13			28	17	18	15	
14			30	21	20	15	
15			31	21	18	16	
16			31	21	18	17	
17			6J	20	19	17	
18			20	20	19	14	
19			27	19	- 17	18	
				19	19	18	Leaflen
Average	- 1	21		19	16	15	
Standard Dev	1.215431087	2.79067712	3.670651742	1.801558779	3.300318964	2.010207868	
T-Test	0.925301893		4.42705E-08		0.638773373		

				Cyclone	472	508	444	479	462	502	461	576	458	432	468	454	476.3	3.81E+01	
Color	7.5gy4/4	5gy 4/8	Head wt. (g)	Darkland Cos Cy	630	299	548	704	453	222	465	455	460	447	340	442	515.7	1.09E+02	2.50E-01
		15			32	36	35	36	35	34	35	98	35	34	35	32	34.8	1.11E+00	
Days to Maturity		1 68	Frame diam (cm	Darkland Cos Cyclone	68 26	38	37	68 36	37	3 40	98 39	37	98	68 39	1 39	37	1 38.0	$1.21E\pm00$	1.03E:06
Date Mature	11/2/01	10/31/01	ore Length	Cyclone	7.7	9:9	6'9	7.5	6.1	6.3	6.8	7.2	9.9	8.6	7.1	7.	7.1	6.79E-01	
	Cyclone	Darkland cos	Head length: Core Length	Darkland Cos Cyclone	7.0	6.4	6.7	7.6	7.3	8.3	5.9	7.1	6.7	8.9	7.0	8.3	7.1	7.00E-01	8.51E-01
			angth (mm)		270	270	270	270	270	270	087	280	270	250	290	280	272.5	9.65E+00	
10/30/01	AC/DG		Head 1	Darkland OCyclone	210 280	210 290	220 290	230 320	210 330	230 290	200 290	220 250	230 260	230 320	240 280	220 290	220.8 290.8	01 2.35E+01	CO2H20/C
Date evald:	10 Eval by:		mm)	Cyclone	240 2	230 2	230 2	270 2	280 2	240 2	240 2	200 2	210 2	220 2	230 2	230 2		1.16E+01	12
Tury			Heart Length (mm)	Darkland Cos		2.							7				235.0	2.24E+01	6.45E-02
Ranch/lot:	ommercial Va		m)	yelone	35	41	39	36	44	43	41	39	41	67	41	39	39.0	4.07E+00	
Salinas	Costa/Mann Commerc		Core Length (mm)	Darkland Cos Cyclone	40	45	43	42	45	35	64	SE	68	47	9	38	41.3	4.77E+00	2.27R-01
Location:	8/24/01 Grower:		(unu)	Суслопе	34	35	36	37	38	38	39	37	35	36	35	34	36.2	1.64E+00	
DS01038	8/24/01		Core Diameter (mm)	Darkland Cos	39	41	32	38	31	35	31	32	34	36	37	98	35.2	3.27E+00	3.54E-01
Trial map #: PDS01038	Wet Date:			Sample #	П	2	3	4	3	9	7	8	6	10	11	12	Average	Stan dev	Ttest

hlighted in yellow indicates a statistically significant difference at a minimum 95% confidence level

2 2 3 3 4 4 4 6 6 6 6 6 6 6 6 6 6 6 6 6 6 6		Darkland Cos 19 20 20 21 17 17 19 10 10 10 10 10 10 10 10 10 10 10 10 10	Cyclone 16 19 19 19 21 21 21 21 21 21 21 21 21 21 21 21 21	Darkland Cos Cyclome 20 21 21 24 25 25 26 26 27 27 28	12 12 12 13 14 14 14 14	Darkland Cos 9 9 12 12 14 14 15 15 15 15 15 15 15 15 15 15 15 15 15	Data highlighted in yellow indicates a statistically significant difference at a minimum Statistically Distinct Confidence Trait Level % Head length 98 Frame diam 99 Leaf Longth 99	confidence Confidence Level % 98 99 99
8 0 0 11 12 12 12 12 12 12 12 12 12 12 12 12	20 20 20 19 19	20 16 16 17 17 18						
17 18 19 Average Average T.Test 0.1	153741223 0.171295933	18	19 20 17 19 1663134193 3.88987E-08	24 26 27 27 3166897499	17 16 15 16 16 2.064741605 0.683826647	19 19 17 16 3,314861097	Loaf length and leaf width numbors are measurements of the useable leafs from one plant.	neasurements of the

					Cyclone		84/	883	4100	111
	7 Sov4/6	5gy4/6		Head wt. (g)	arkland		1,11,1	822	3701	1003
ſ	Ť	5		H	Cvolone	70	ŧ.	37	200	34
Days to Medicites	86	92		rame diam (cm	Darkland	41	-	4	ç	7
Date Mature	5/13/02	20/1/92		undirem a	yclone	4.5		, 0,0	46	O'F
	Cyclone	Darkland	E POST STORY	cosmental man	Darkland	5.4		5,5	0.9	25
				(mm) ms	Cyolone	027		007	280	
5/7/02	dg/ac		New Allers	Tormour .	Larkland	270		Z/#U	300	
Date evald:	Eval by:				Cyclone	210	300		220	
Salmina 2	I Var: El Camino Real Eval by:		Heart Conorth Car	D-11.4	DISTRICT	230	180		260	220
Sanch/lot:	Commercial Var:		(44)	Yerlone	A CONTRACTOR CONTRACTO	9	\$		19	02
RSV02026 Location: Soledad Ranch/lot:	Costa/Mann Commercial		Core Length (n	Darkland		39 50	5 45		2 20	40
Location:	2/4/02 Grower:		ter (mm)	Cyclone			7		22	37
RSV02026	2/4/0		Core Diameter (Darkland	ľ	30	2 37	,	S.	4
#	Wet Date:			Sample #	ļ					-

	2/4/02	2/4/02 Grower:	Costs/Afann	Costs Mann Commercial Vom	DI Commission Days 1						Call to trade the	arus rey	CUIUT	
			-	Commercial var.	П		agy ac		Cyclone	5/13/02	86		7.5gv4/6	
									Darkland	2/1/02	92		5gy4/6	
	Core Diameter (mm)	r (mm)	Ches T month		e de la constante de la consta									
			run) manor o no	from	THEATT PORRU (MM)	Ú	Head Jength (mm)	ch (mm)	Head length: Core Length		Frame diam (cm)		Head wt. (2)	
	Darkland	Cyclone	Darkland	Cyclone	Darkland	Cyclone	Darkland	100 miles	Darkland	1000000	Doeblend	Total Actor	Г	
1	36	39	05 6	09	230		4	270		Amounto	1		Darkiand	Cyclone
124	57	\$5	45					017			41		744	847
ľ				*		7007	740	250	5.3	5.0	4	37	822	883
"	52	52	20	0 61	1 260	220	300	280	6.0	4.6	42		1064	444
4	55	37	7	09	0 220	190	300	270	0.9		Ę		100	///
4)	52	39	55	19	230	2000	02.0	OFC			1		76/	993
ľ	96						2/7	210	4.9	4.4	42	34	955	945
"	9		20	0 62	250	220	300	260	0.9	4.2	40	35	1120	707
`	43	37	40	0	5 210	210	300	250	7.5		200	0.0		127
00	42	30	9	95	000	901	o Lo				7	ΛC	/41	10/
ľ						130	0/7	0007	5.4	4.5	4	39	820	957
^ ;		39	8	55	5 230	200	300	260	0'9	4.7	39	300	810	680
2		41	45	5 60	210	210	300	280	6.7	4.7	41	36	1175	1134
티	55	36	50	0 65	5 220	210	300	260	6.0		QP	3.0	200	7001
12	39	55	5 45	5 62	200	220	280	270	6.2		3.8	35	C76	1050
	47.8	43.4	1 48.3	3 59.7	7 221.7	206.7	285.8	264.2			707	6	,0,0	100
	7.25E+00	7.61E+00	3.89E+00	0 4.36E+00	2.12E+01	1.07E+01	1 085-01	1.0812-01	302 9	2.1.5	40	,	C.00%	1.01.6
	1676-01		CA CT2 00				10.000	TA OTALA	10-22/120	3.175.01	1.10E:T0V	7.04E+00	1.35E+02	1.14E+02
	1007/001		V.30E-V	,,	4.00E-02		3.03E-03		5,41E-07		4.94E-06		9.03E-01	

Indicates that the varieties are significantly different for this trail at least a 93% confidence level

	Confidence	Level %	66	96	66	66	66	66	66
	Statistically Distinct	Trait	Core Length	Heart Length	Head length	Head length:Core Length	Frame diam	LeafLength	Leaf Thickness
•									

	Leaf County Plant	lant	Leat Length	(MCC)	Leat Width	(mm)	Leaf Thickner (mm)	(um)
	Darkland	Cyclone	Darkland	Сусіопе	Darkland	Cyclone	Darkland	Cyclone
	35	35	797	7 231	144	141	96'0	0.95
2		32	277	192	182	126		
ඟ ි	37	28	760	177	127			26'0
4	37	28	270	245	142	154		560
5	35	78	282	861 2	158	110		660
9	34	39	092	0 233	188	182		6.63
7	33	37	261	218	171			\$60
∞	28	35	270	722	111			86'0
6	36	34	277		155	180		96'0
10	36	34	240	243	162	155	0.81	0.82
11	35	32	251	248	167	180	0.84	86'0
12	33	32	283	254	113	187		60
13			280	262	153	1771	0.84	0.82
14			275	218	157	141	98'0	0.82
15			265	5.27	157	152		60
16			258	339	175	180		760
17			255	252	203	178	160	(4)
18			240	248	195	159	0,92	1,29
19			270	238	215	162	0.85	960
20			275	233	205	160	0.85	16'0
Average	34	33	266	234	164	851	98'0	9660
tandard Dev	2.429303429	3.84550111	Ø.	17,79554941	28.87905816	20.94252787	0,04518966	0.158426475
T-Test	1.96E-01		111111111111111111111111111111111111111	1/	4.42R-01		120,201	

				_						
					Cembono	Cyclotte	22,6	6/6	953	1120
	Color	7 Smd/4	/2,6/6 5gy5/6	77.7	Fload Wt. (g)	000	1100	1110	9111	1530
				Salara Constitution (Access	Cologe	30	66	300	00	t or
	ays to Maturity	80		The second	Darkland	, i	000	8 6	000	8
	Date Mature	6/7/02	6/3/02	a I enoth	Cyclone	9.5	2.0	4.0	64	43
	<u> </u>	Cvolone	Darkland	Head landh Over I south	Darkland C	4.2	40	0.0	6.0	4.4
				eth (mm)	Cyclone	237		066		252
	5/31/02	AC		Head let	Darkland	250		2		292
	Date evald:	Eval by:		mm)	Cyclone	154 165		69 162	84 182	179 179
0,0	1219	Frontier		Heart Length (1	Darkland	15	170	16	18	17
	Kanch/lot:	Commercial Va Frontier		m)	Cyclone	42	59	47	57	59
	1	D'arrigo (Core Length (mm	Darkland	49	49	63	57	19
T a a a different	Location:	3/10/02 Grower:		r (rom)	Cyclone	40	36	41	37	37
Detrocan	2	3/10/02		Core Diameter (mm	Darkland	42	38	34	45	44
'rial mon #	إ	Vet Date:			ample #	1	2	3	4	\$

			_				سيت												
				Curlond	O O	070	910	1130	27.8	808	1040	786	\$00	1013	1156	1286	9046	1 44R+02	
700	5010r	/.Jgy4/+ 5gy5/6		Head Wt. (g)	1060	1186	1110	1350	1539	855	928	1031	823	1005	1420	1270	1132.2	2.26F±02	8.97E-02
		5		olyan	3.5	65	32	96	55	32	32	34	38	33	45	34	33.7	1.67F+00	
ove to Medumite	20 1	85	STATE STATE OF THE	Parkiand Co	2 30	86	128	98	39	34	37	37	37	37	38	36	36.8	1.27E+00	3.00E-05
Date Mature lo	- 2	6/3/02			5.6	44	4.0	4.2	4.3	4.6	4.2	4.2	6,4	5.4	49	4.7	4.8	6.83E-01	
4	volone	Darkland	Head landby Cost Land	Darkland Cv	5.3	4.9	3.9	5.0	4.4	5.4	5.2	0.9	4.8	5.8	6.5	5.3	5.2	7.18E-01	1.91E-01
L	Ć	iä			237	262	229	239	252	269	262	239	249	237	224	205	242.0	1.81E+01	
5/31/02	AC	7	Flast leneth (mm)	Darkinnd	259	239	244	283	292	292	253	259	366	283	320	287	273.1	237E+01	1.54E-03
Date evald:	ł			yclone	165	190	162	182	179	192	205	179	172	159	166	169	176.7	1.39氏+01	
1219	Va Frontier		Heart Length (mm)	Darkland	154	170	169	184	179	184	159	157	169	167	172	159	168.6	1.01E+01	1.17压-01
Ranch/lot: 12	Commercial Va Fr			lone	42	59	47	57	59	59	62	57	39	44	46	44	51.3	8.25E+00	
Salinas Rat	D'arrigo Cot		Core Length (mm)	Darkland Cyclone	49	49	63	57	19	54	49	43	26	49	49	54	53.3	6.76E+00	5.23E-01
			Γ	Cyclone Da	40	36	41	37	37	36	35	40	32	45	40	42	38.4	3.55E+00	
RSV02049 Location:	3/10/02 Grower:		Core Diameter (mm)	Darkland C	42	38	34	45	4	42	39	39	40	38	40	42	40.3	2.99E+00	1.85E-01
	1 1		_		-		_	_	-	_			-	_	_	_	_		_

650	
1453	
ily different for this trait	
520	
623	MON.
86-01	1.
1990	
different for th	
1,070	1000
120	100
15.3	1.0
1	100
100	1.128
10027	1000
(2)	100
27	1567
1/4	10000
16	122.5
25.7	A 200
200	100
100	490
102	1000
: 22	03.50
2.70	
(S)	155
765	315
123	
100	100
18.55	/ S. C.
100	0.37230
2000	100
12.00	100
45.5	1.0
16.00	- 40
200	AN 9 📆
833	116
127	100
12.0	1.5
177	Z > 1
100	A
230	111
	. 12
16 E	2.0
(2)	0.00
577	120
es that the varieties are gignificantly	
100	333 C
15343	14907
10.2	20.00
100	>
1.4	
2000	69 D
12.04	
120	
177	
1.0	4.40
1,74	1.74
12000	
160,000	
200	100
15.00	2.70
197	2.0
723	
# 0	
(a = 0	100
100	100
- ≪≃	in.
	2.34
(1 -1	0.00
Indicate	at at least a 95% confidence level

•			
	Statistically Distinct	Confidence	
	Trait	Level %	
	Jead length	98.5	
	rame diam	86	
	Head Weight	91	
	Leaf Length	66	
	eaf Thickness	5'96	

Leaf Thickness	Darkland Cyclone	0.70 0.80	0.70	0.80	080 0.60					0.50 0.70	0,40 0,50	0.50 0.70	0.50 0.50	0.50 0.60	0.40 0.50			0.50	0,40 0,60	0.80 0.80	0.70 0.80	1 0,630	Carried Control of the Control of th
当	Cyclone D	168	192	125	186	177	146	152	14	164	168	136	189	186	165	139	142	160	169	163	159	162	3 - to otto to to o t
Leaf Width	Darkland	171	158	152	160	192	171	168	138	152	157	178	174	154	149	146	139	148	175	175	159	191	14 32 400010
	Cyolone	227	214	220	222	219	235	243	234	238	232	215	204	223	219	225	214	239	209	228	238	225	UUU22 saa UE
Leaf Length	8000	246	210	997	692	797	252	233	246	220	752	742	277	240	283	279	233	275	229	219	245	250	35300005 50
lant	Cyclone	24	31	36	31	32	72	35		62	98	67	33									31	Standard Day 2 014112446 2 77602240
Leaf Count/ Plant	Darkland	41	36	37	36	32	38	98	36	38	29	37	98									36	3112112
	Sample #	1	2	3.	4	5	9	7	8	6	10	11	12	13	14	15	16	17	18	19	20	Average	Standard Day

Whole Leaf Process Data

Data produced by Fresh Leaf Farms, LLC in conjunction with Mann Packing

9	Frontier	1.11	1.05	1.18	0.97	0.93	1.048	1.02E-01	
Pack Out %	Fro	191	1.1	44.	1.18	1.03	27	ـــ	22
Pac	Cyclone	Ţ			,		1.272	2.44E-01	4.63E-02
SU	Frontier	616	009	962	839	1029	8.698	1.66E+02	
Total Cartons	Cyclone	630	781	955	1001	1121	915.6	2.09E+02	7.11E-01
	Frontier	828	569	812	898	1104	836.2	1.90E+02	
Trays Processed	Cyclone	392	707	999	924	1092	755.4	2.67E+02	5.96E-01
Process	Date	15-May	3-Aug	22-Aug	27-Aug	23-Sep			

10 pounds of processed and stacked whole leaves of romaine, all of which meet a specific size requirement. a single tray of product. All trays contain 24 harvested romain plants. All finished cartons are This data compares the pack out proportion of the lettuce varieties Cyclone and Frontier. The pack out number is the proportion of total cartons (finished product) generated from

For example, based on the May 15 data:

For Cyclone, 1.61 finished 10 pound cartons were produced from a single tray of 24 romaine plants. For Frontier, 1.11 finished 10 pound cartons were produced from a single tray fo 24 romaine plants. Based on the 5 compared dates, Cyclone is statistically higher yielding in this patented process at a 95% confidence level.



Definition of the plant characteristics as measured in our trials:

- 1. The romaine plant is cut according to fresh market standards, at market maturity. The plant is weighed, the core diameter is measured, and then the plant is sliced vertically.
- 2. We define the heart as the part of the plant enclosed by the outer most cupping leaf. This area is the densest part of the plant and the color varies from white to yellow to light green as the leafs mature.
- 3. For our evaluation purposes the heart length is measured from the base of the cut plant stem (core) to the tip of the outer most cupping leaf that forms the heart.
- 4. The stem length is measured from the base of the cut stem to the stem tip, at market maturity.
- 5. Head height is measured vertically from the base of the cut stem to the tip of the margin of the longest leaf.
- 6. Head height differs from plant height. Plant height would be the total plant height of the growing plant from the top of the bed to the highest part of the plant. Head height is a measurement of the plant height once cut to market standards.





Trialing Protocol for Intellectual Property Protection.

I. Set Up

- 1. Parental lines and competing varieties are identified.
- 2. Primary slots are identified.
- 3. Necessary accession lines are located and purchased/received from seed dealers or growers.
- 4. All varieties are assigned a number to maintain integrity, and anonymity.
- 5. Trials are set up in the Progeny warehouse with all necessary varieties. Variety arrangement for trial is diagramed.

II. Planting

- 1. Commercial plantings are located by contacting commercial growers during the planting slot recommended for the variety.
- 2. Field is located during commercial planting, and the necessary rows and area is marked off by a Progeny employee with proper training.
- 3. Varieties are planted according to diagram, in 100 ft. ranges.
- 4. All varieties are planted in same manner, to mimic the planting of the commercial variety as closely as possible.
- 5. A trial map is drawn diagramming the trial, the trial location in the field, and directions to the field.

III Maintenance

- 1. All varieties are treated identically. The grower handles all watering, fertilization, and pest control, as if it was no different from the commercial field it is grown with.
- 2. Thinning of the trial is done by a crew contracted by the commercial grower.



IV. Evaluation

- 1. Evaluations are done as near to the time of the commercial harvest as possible by knowledgeable Progeny employees.
- 2. The evaluation is conducted "blindly". The evaluator(s) do not have the key to the trial at the time of evaluation.
- 3. 24 heads of each variety are evaluated.
 - a. The frame diameter of 12 random plants are measured to the nearest cm.
 - b. 12 mature heads of each variety are cut to market specs.
 - c. The heads are carried to an adequate work station
 - d. The following measurements are then conducted and recorded:
 - 1. Each head is weighed to the nearest gram.
 - 2. The core diameter of each head is measured to the nearest mm.
 - 3. The heads are then sliced in to halves, discarding 1 half.
 - 4. The core lengths (from the cut stem to the core tip) are measured to the nearest mm.
 - 5. The heart lengths (from the cut stem to the tip of the longest heart leaf) are measured to the nearest mm.
 - 6. The head length (from the cut stem to the tip of the longest leaf) is measured to the nearest mm.
 - 7. The ideal maturity or harvest date is then estimated based on the solidity of the head, the core length and any other physiological characteristics present.
 - 8. The leaf color is documented using the Munsell Color Charts for Plant Tissue.
 - e. From these measurements, we then use an Excell program to calculate the averages, the standard deviations and the T-Tests for the compared varieties.

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE	The following statements are made in	n accordance with the Privacy Act of
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is required in order to decertificate is to be issued (7 U.S.C. 2 until certificate is issued (7 U.S.C. 2426	rk Reduction Act (PRA) of 1995. eterminé if a plant variety protection 4211 - Information is held as 5
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION	3. VARIETY NAME
PROGENY ADVANCED GENETICS, INC.	OR EXPERIMENTAL NUMBER PX 408	CYCLONE
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (include area code)	6. FAX (include area code)
1536 B MOFFETT ST	831-751-6030	831-751-6032
SALINAS CA 93905	7. PVPO NUMBER	
	200	300054
8. Does the applicant own all rights to the variety? Mark an "X" in appropri	nate block. If no, please explain.	YES NO
	•	,
is the applicant (individual or company) a U.S. national or U.S. based or If no, give name of country	ompany?	(YES NO
10 is the applicant the original general	() If no, please answer one of the fo	
a. If original rights to variety were owned by individual(s), is (are) the or YES YES b. If original rights to variety were owned by a company(ies), is(are) the	IO If no, give name of country	?
☐ YES ☐ N	IO If no, give name of country	
11. Additional explanation on ownership (if needed, use reverse for extra sp	pace):	
		:
	,	
PLEASE NOTE:		
Plant variety protection can be afforded only to owners (not licensees) who meet on	e of the following criteria:	
 If the rights to the variety are owned by the original breeder, that person must be which affords similar protection to nationals of the U.S. for the same genus and s 	a U.S. national, national of a UPOV member	
 If the rights to the variety are owned by the company which employed the origin member country, or owned by nationals of a country which affords similar protect 	al breeder(s), the company must be U.S. base tion to nationals of the U.S. for the same ge	nus and species.
3. If the applicant is an owner who is not the original owner, both the original owner		
The original breeder/owner may be the individual or company who directed final bree		

According to the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to compete this information collection is estimated to average 10 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. The U.S. Department of Agniculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and manual or familial status. (Not all propriate bases apply to all programs). Persons with disabilities who require atternative means for communication of program information (braille, large print, audiotape, etc.) should contact

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal

STD-470-E (07-97) (Destroy previous editions).

3/